

Microcystin LF


from *Microcystis aeruginosa*



Analytical Standard

Product Information

Cat. No.	Amount
MC-LF-a	1 mL (about 10 µg/mL)



Product Specifications

Molecular Formula	C ₅₂ H ₇₁ N ₇ O ₁₂
Molecular Weight	985.5 g/mol
Purity	>95 % (HPLC)
Source	<i>M. aeruginosa</i> strain
Form	solution of about 10 µg/mL in methanol
Shipping	Ambient
Long Term Storage	- 20°C
Shelf life	24 months
Stability	The analytical standard should be used immediately after the vial is opened

Description

Cyclic heptapeptide toxin isolated from the freshwater cyanobacterium *Microcystis aeruginosa*.¹

The identity of the compound has been confirmed by NMR and MS/MS.

The analytical standard is dissolved in 100% methanol and ready to use for calibration. It is distributed in amber glass vials containing around 10 µg in 1 ml MeOH. The concentration of each lot is determined spectrophotometrically, confirmed by HPLC, and stated on the Certificate of Analysis.

For research use only!

Not available for sale to end-users without signing an end-use-certificate as required by German and international law.

[1] Azevedo et al., First report of microcystins from a Brazilian isolate of the cyanobacterium *Microcystis aeruginosa*, *J Appl Phycol.* 1994, 261-265

Lawton et al., Isolation and characterization of microcystins from laboratory cultures and environmental samples of *Microcystis aeruginosa* and from an associated animal toxicosis, *Natural Toxins.* 1995, 50-57